arXiv:1502.02858v1 [gr-qc] 10 Feb 2015

On the local form of static plane symmetric space-times in the presence of matter

L. G. Gomes

Instituto de Matemática e Computação
Universidade Federal de Itajubá , Brazil

Abstract

For any configuration of a static plane-symmetric distribution of matter along space-time, there are coordinates where the metric can be put explicitly as a functional of the energy density and pressures. It satisfies Einstein equations as far as we require the conservation of the energy-momentum tensor, which is the single ODE for self-gravitating hydrostatic equilibrium. As a direct application, a general solution is given when the pressures are linearly related to the energy density, recovering, as special cases, most of known solutions of static plane-symmetric Einstein equations.

1 Introduction

The gravitational field in the vicinity of a large and isolated distribution of matter is approximately static if the interval of time considered is small compared to the characteristic time the system changes. As the matter distribution does not suffer any substantial change except in only one direction, it is also plane symmetric. This is the case, for example, when we are dealing with Newtonian gravitation close to the surface of a planet. In this case, define the height from the ground as z, the gravitational potential as ϕ and the atmospheric mass distribution and pressure as ρ and p, respectively. They are approximately functions of z only. Besides an appropriate equation of state relating the later thermodynamic variables, there is the hydrostatic equilibrium [1]

$$\frac{dp}{dz} = -\rho \, \frac{d\phi}{dz} \qquad , \tag{1}$$

with the Newtonian potential a solution of Poisson equation $\frac{d^2\phi}{dz^2} = \frac{1}{2}\,\rho$. Here units are such that $8\pi G = 1$ and $c = 1$.

The gravitational equation is simple enough to be integrated and give ϕ as

$$\phi(z) = g\, z + \frac{1}{2}\int_0^z du \int_0^u dw\, \rho(w) \qquad , \tag{2}$$

where g is the constant acceleration of gravity in the absence of atmosphere. Using eq.(1) and eq.(2) we deduce the self-gravitating equation for hydrostatic equilibrium:

$$\frac{d}{dz}\left(\frac{1}{\rho}\frac{dp}{dz}\right) = -\frac{1}{2}\rho \qquad . \tag{3}$$

To set the stage for the relativistic generalization of the situation above, we start with a four dimensional Lorentzian manifold M admitting an isometric action of the transformation group of the Euclidean plane with 2-dimensional space-like orbits. Let e_1 and e_2 stand for the induced killing vector fields corresponding to translations in orthogonal directions. We also assume there is another killing vector field e_0 which is time-like, orthogonal to and commuting with e_1 and e_2. Choosing a curve $\gamma(z)$ such that $\dot{\gamma}(z) \neq 0$ is orthogonal to e_i, $i = 0, 1, 2$, and spreading it all around using the fluxes of the three killing vector fields, we define a coordinate system where $e_0 = \frac{\partial}{\partial t}$, $e_1 = \frac{\partial}{\partial x}$, $e_2 = \frac{\partial}{\partial y}$ and the metric is

$$ds^2 = g_{00}(z)\, dt^2 + g_{11}(z)\,(dx^2 + dy^2) + g_{33}(z)\, dz^2 \quad . \tag{4}$$

We also assume that the energy-momentum tensor is equally symmetric, such that in these coordinates it is diagonal

$$(\, T^{\mu}_{\nu}\,) = \mathbf{diag}\,\{\, \rho(z), -p_{||}(z), -p_{||}(z), -p(z)\,\} \quad . \tag{5}$$

Note that the energy density $\rho = T^0_0$, the pressure parallel to the plane of symmetry $p_{||} = -T^1_1 = -T^2_2$ and the pressure orthogonal to it $p = -T^3_3$ behave as functions under a coordinate change of the type $u = u(z)$. This implies they are all defined globally in such space-time. To simplify our analysis, we set the pressure difference δp as

$$\delta p = p - p_{||} \quad . \tag{6}$$

Such scheme defines what we mean by a static plane symmetric space-time (see also [2], ch. 15).

Unlike Newtonian physics, where the gravitational part is played by one potential and one linear Poisson equation, in the relativistic analogue we deal with three unknown functions playing the hole of the potential, g_{00}, g_{11} and g_{33}, and three non-linear independent Einstein equations, which are far from being easily integrated (see [3], for instance). Indeed, the component g_{33} is spurious as we can make it equal unit in a simple coordinate change $du = g_{33}(z)\, dz$.

Although we may identify matter as a fluid, in the relativistic approach it is natural to consider a more complex and general structure of the energy-momentum tensor. Therefore there is no need to consider an isotropic stress, that is, $\delta p = 0$. As an example, for an electric charged plane [3] we have $\delta p = 2p = -2\rho$. As a result, our complementary equations of state are two in number.[1] This makes our system well defined: we have five independent unknown functions, g_{00}, g_{11}, ρ, p and δp, related by five independent equations: three of them coming from the gravitational part and the rest from the sort of matter present in the system. In this paper, our main result is to give g_{00}, g_{11} and g_{33} in terms of ρ, p and δp such that they satisfy Einstein equations, as we explain bellow.

---

[1] By an equation of state we mean any one coming from the matter source, which could be in the form of an algebraic equation, as it is usual in thermodynamics, as a differential equation coming from a lagragian, etc.

Define the metric

$$ds^2 = e^{2\phi}\, dt^2 - e^{4(z_2-z_1)\int_0^z dw\, \frac{\phi'}{w-z_2}} \left(dx^2+dy^2\right) - \frac{4\,(z_2-z_1)\,(z-z_1)}{(z-z_2)^2}\,\frac{(\phi')^2}{p}\, dz^2 \quad , \tag{7}$$

where $\phi = \ln(g_{00})^{\frac{1}{2}}$ is the generalized gravitational potential given by

$$\phi = \int_0^z dw\, \frac{w-z_2}{w-z_1}\left(\frac{p}{(w-z_1)(\rho-p)+(z_2-z_1)(\rho+7p-4\delta p)}\right) \quad . \tag{8}$$

We also assume that the derivative ϕ' is continuous and different from zero in a neighbourhood of $z=0$. Thus $p \neq 0$ close to $z=0$. The arbitrary constants z_1 and z_2 satisfy $z_2 > z_1$ and $z_1\, p_0 < 0$, as p_0 is the value of p at $z=0$.

The conservation of the energy-momentum tensor (5) in the space-time with the metric above, $\nabla_\mu T^\mu_\nu = 0$, turns out to be the generalization of the hydrostatic equilibrium relation

$$\frac{dp}{dz} = -\left(\rho + p + \frac{4\,(z_2-z_1)}{z-z_2}\,\delta p\right)\frac{d\phi}{dz} \quad . \tag{9}$$

This is commonly find in the literature for an unknown ϕ and $\delta p = 0$ ([4], sec. 5.4). If we apply the definition of ϕ (eq.(8)), we arrive at the relativistic self-gravitating hydrostatic equilibrium equation

$$\frac{dp}{dz} = -\left(\frac{(z-z_2)\,(\rho+p) + 4\,(z_2-z_1)\,\delta p}{(z-z_1)\,(\rho-p) + \,(z_2-z_1)\,(\rho+7p-4\delta p)}\right)\frac{p}{z-z_1} \quad . \tag{10}$$

In this paper we show that the metric in eq.(7) and the energy-momentum tensor in eq.(5) satisfy Einstein equations , that is, $R^\mu_\nu - \frac{1}{2}R\,\delta^\mu_\nu = T^\mu_\nu$, provided the conservation above is attained. Furthermore, the case when ϕ cannot be defined as in eq.(8) is also explained. These are presented in section 2. In the following, we show how it simplifies the analysis of space-times in the presence of fluids with the prescribed equations of state $p = (\gamma(\rho)-1)\,\rho$ and $\delta p = \epsilon(\rho)\,\rho$, giving in a straightforward and unified way most of the solutions studied so far, which fall in the category of γ and ϵ constants. In the last section we make our final remarks.

2 The local form of the metric

Our main result is given in the following theorem:

Theorem 1 *The metric given in eq.(7) and the energy-momentum tensor in eq.(5) satisfy Einstein equations in a neighbourhood of $z = 0$, provided the later attains the self-gravitating hydrostatic equilibrium given in eq.(10).*

Proof: We proceed straightforwardly in computing the Einstein tensor for the metric (7). Thus, the non-vanishing components of the Levi-Civita connection are

$$\Gamma^t_{tz} = \phi' \quad ; \quad \Gamma^x_{xz} = \Gamma^y_{yz} = 2\,\frac{(z_2 - z_1)\,\phi'}{z - z_2} \quad ; \quad \Gamma^z_{tt} = \frac{e^{2\phi}\,(z - z_2)^2\,p}{4\,(z - z_1)\,(z_2 - z_1)\,\phi'}$$

$$\Gamma^z_{xx} = \Gamma^z_{yy} = -\frac{(z - z_2)\,p}{2\,(z - z_1)\,\phi'}\; e^{4(z_2 - z_1)\,\int_0^z dw\,\frac{\phi'}{w - z_2}} \quad ; \quad \Gamma^z_{zz} = \frac{1}{2}\,\frac{d}{dz}\,\ln\left(\frac{4\,(z_2 - z_1)\,(z - z_1)}{(z - z_2)^2}\,\frac{(\phi')^2}{p}\right)$$

Through the rest of our proof we assume $\nabla_\mu\, T^\mu_\nu = 0$, which is equivalent to eq.(9). Therefore, during the calculation of the curvature $R^{\lambda\mu}_{\kappa\nu} = g^{\mu\mu'}\,(\partial_\kappa\,\Gamma^\lambda_{\nu\mu'} - \partial_\nu\,\Gamma^\lambda_{\kappa\mu'} + \ldots)$ we apply the identities

$$\frac{p'}{\phi'} = -\left(\rho + p + \frac{4\,(z_2 - z_1)}{z - z_2}\,\delta p\right) \tag{11}$$

and

$$\frac{p}{\phi'} = \frac{z - z_1}{z - z_2}\,\left((z - z_1)(\rho - p) + (z_2 - z_1)(\rho + 7p - 4\delta p)\right) \quad , \tag{12}$$

the last one coming from the very definition of ϕ. Then we obtain for its independent components

$$R^{tx}_{tx} = R^{ty}_{ty} = \frac{(z - z_2)\,p}{2(z - z_1)} \quad ; \quad R^{tz}_{tz} = \frac{1}{2}\,(\rho + p - 2\,\delta p) + \frac{z_2 - z_1}{z - z_1}\,p \tag{13}$$

$$R^{xy}_{xy} = \frac{z_2 - z_1}{z - z_1}\,p \quad ; \quad R^{xz}_{xz} = R^{yz}_{yz} = -\frac{1}{2}\left(\rho + \frac{z_2 - z_1}{z - z_1}\,p\right) \quad . \tag{14}$$

For the Ricci tensor $R^\mu_\nu = R^{\lambda\mu}_{\lambda\nu}$ and scalar curvature we have

$$R^t_t = \frac{1}{2}\,(\rho + 3\,p - 2\,\delta p) \qquad ; \quad R^x_x = R^y_y = -\frac{1}{2}\,(\rho - p)$$
$$R^z_z = -\frac{1}{2}\,(\rho - p + 2\,\delta p) \quad ; \quad R = -\rho + 3\,p - 2\,\delta p \quad .$$

Hence Einstein equations hold for ds^2, as can be readily verified.

□

It is not true that ds^2 in eq.(7) can be well defined for any kind of matter. In order to be complete, the following theorem explains when such a scheme is not possible.

Theorem 2 *If ϕ' is nowhere well defined or if it vanishes everywhere, then around each point of this static plane symmetric space-time one of the following holds, with ds^2 satisfying Einstein equations:*

(i) $p=\rho=0\ \ ,\ \delta p=\delta p(z)$. *Defining* $\varphi=-\int_0^z dw\,\frac{w-z_1}{\delta p+(w-z_1)^2}$ *, we have*

$$ds^2=e^{2\,\varphi}\,dt^2-\left(dx^2+dy^2\right)-\left(\frac{\varphi'\,dz}{z-z_1}\right)^2 \quad . \tag{15}$$

(ii) $p=0\ \ ,\ \rho=4\,\delta p\ \ ,\ \delta p=\delta p(z)$. *Defining* $\varphi=\frac{1}{3}\int_0^z dw\,\frac{w-z_1}{3\,\delta p+(w-z_1)^2}$ *, we have*

$$ds^2=e^{2\,\varphi}\,dt^2-\,e^{-\,4\,\varphi}\left(dx^2+dy^2\right)-\left(\frac{3\,\varphi'\,dz}{z-z_1}\right)^2 \quad . \tag{16}$$

(iii) $\rho=p\,,\,\delta p=2\,p$. *There are constants* α *and* β *such that*

$$p=\frac{4\,\alpha\,\beta}{3\,(1+(\,\alpha+\beta\,)\,z\,)^2} \quad . \tag{17}$$

If $\alpha+\beta\neq 0$,

$$ds^2=(1+(\,\alpha+\beta\,)\,z\,)^{\frac{2}{3}\left(\frac{3\alpha-\beta}{\alpha+\beta}\right)}\,dt^2-\,(1+(\,\alpha+\beta\,)\,z\,)^{\frac{4}{3}\left(\frac{\beta}{\alpha+\beta}\right)}\left(dx^2+dy^2\right)-\,dz^2 \quad , \tag{18}$$

and if $\alpha+\beta=0$,

$$ds^2=e^{\frac{8}{3}\,\alpha\,z}\,dt^2-\,e^{-\,\frac{4}{3}\,\alpha\,z}\left(dx^2+dy^2\right)-\,dz^2 \quad . \tag{19}$$

In the special case $p=0$ *we obtain, for* $\beta=0$*, the Minkowski metric described by an observer with a uniform acceleration* α *([5]) or, for* $\alpha=0$*, the Taub-Levi-Civita vacuum solution ([3]) .*

Proof: If we take the metric (4) in coordinates such that $du=\sqrt{-g_{33}}\,dz$ and define[2]

$$\xi(u)=\frac{3}{4}\,\frac{d}{du}\,\ln|\,g_{11}\,| \qquad \text{and} \qquad \psi(u)=\frac{d}{du}\left(\frac{1}{2}\,\ln g_{00}+\frac{1}{4}\,\ln|\,g_{11}\,|\right) \quad , \tag{20}$$

such that

$$ds^2=e^{\frac{2}{3}\int du\,(3\psi(u)-\xi(u))}\,dt^2-\,e^{\frac{4}{3}\int\,du\,\xi(u)}\left(dx^2+dy^2\right)-\,du^2 \quad , \tag{21}$$

then Einstein Equations turn into

$$G^t_t=T^t_t \quad : \quad -\frac{4}{3}\left(\xi'+\xi^2\right)=\rho \tag{22}$$

$$G^t_t-4G^x_x=T^t_t-4T^x_x \quad : \quad 4\left(\psi'+\psi^2\right)=\rho+4p-4\delta p \tag{23}$$

$$G^u_u=T^u_u \quad : \quad -\frac{4}{3}\,\xi\,\psi=-p \quad . \tag{24}$$

---

[2] This is a further simplification for the form of the metric appearing in [6].

If the derivative of the potential is zero everywhere then $p = 0$. Thus, from eq.(24), we conclude that $\xi = 0$ or $\psi = 0$.

If $p = 0$ and $\xi = 0$, then $\rho = 0$ as eq.(22) demands. Defining the coordinate function $z = \psi + z_1$, for an arbitrary constant $z_1 \neq \sqrt{-\delta p(0)}$, from eq.(23) we obtain

$$dz = -\left(\delta p + (z - z_1)^2 \right) du \quad , \tag{25}$$

implying formula (15). This proves (i).

If $p = 0$ and $\psi = 0$, then $\rho = 4\,\delta p$ follows from eq.(23). As before, defining the coordinate function $z = \xi + z_1$ and using eq.(22) we get eq.(16), hence proving (ii).

If the derivative of the potential is nowhere defined then $\rho = p$ and $\delta p = 2\,p$. From eq.(22)-(24), we obtain the following system of ODE's:

$$\xi' + \xi^2 + \psi\,\xi = 0 \qquad \psi' + \psi^2 + \psi\,\xi = 0 \quad . \tag{26}$$

Its general solution is, after we put $z = u$,

$$\psi = \frac{\alpha}{1 + (\alpha + \beta)\,z} \qquad \xi = \frac{\beta}{1 + (\alpha + \beta)\,z} \quad . \tag{27}$$

Returning this expressions in the previous formulas, we prove (iii). □

3 Applications to simple fluids

To illustrate the theorems above, assume we have equations of state in the form

$$p = (\gamma(\rho) - 1)\,\rho \qquad \text{and} \qquad \delta p = \epsilon(\rho)\,\rho \quad . \tag{28}$$

In this case, the configuration of the system is completely determined solving the simple first order ODE of hydrostatic equilibrium for $\rho(z)$:

$$\left(\frac{d\gamma}{d\rho}\,\rho + \gamma - 1 \right) \frac{d\rho}{dz} = -\left(\frac{(z - z_1)\,\gamma + \,(z_2 - z_1)\,(4\epsilon - \gamma)}{(z - z_1)\,(2 - \gamma) + \,(z_2 - z_1)\,(7\gamma - 4\epsilon - 6)} \right) \frac{(\gamma - 1)\,\rho}{z - z_1} \quad . \tag{29}$$

Even though this ODE is not exactly solvable, any approximation method for its solution reflects instantaneously in the metric components, as demanded by equations (7), (8) and (28). This is a great simplification in the analysis of such systems.

Specializing to constant γ and ϵ, such that $\gamma \neq 1$, $\gamma \neq 2$ and $7\,\gamma \neq 4\,\epsilon + 6$, eq.(29) is easily integrated as

$$\frac{\rho}{\rho_0} = \left(1 - \frac{z}{z_1} \right)^{\frac{\gamma - 4\,\epsilon}{7\,\gamma - 4\,\epsilon - 6}} \left(1 - \frac{z}{\alpha\,z_1} \right)^{\frac{-6\,\gamma^2 + 4\,\gamma + 8\,\epsilon}{(2 - \gamma)(7\,\gamma - 4\,\epsilon - 6)}} \quad , \tag{30}$$

where $\rho_0 = \rho(0)$ and

$$\alpha = 1 - \left(1 - \frac{z_2}{z_1}\right) \left(\frac{7\,\gamma - 4\,\epsilon - 6}{\gamma - 2}\right) \quad . \tag{31}$$

Applying it in eq.(7), we arrive to the expression for the metric

$$\begin{aligned} ds^2 = &\left(1 - \tfrac{z}{z_1}\right)^{\frac{2\,(1-\gamma)}{7\,\gamma-4\,\epsilon-6}} \left(1 - \tfrac{z}{\alpha\, z_1}\right)^{-\frac{2\,(1-\gamma)(6\,\gamma-4\,\epsilon-4)}{(2-\gamma)(7\,\gamma-4\,\epsilon-6)}} \; dt^2 \\ &- \left(1 - \tfrac{z}{z_1}\right)^{-\frac{4(1-\gamma)}{7\,\gamma-4\,\epsilon-6}} \left(1 - \tfrac{z}{\alpha\, z_1}\right)^{\frac{4(1-\gamma)}{7\,\gamma-4\,\epsilon-6}} \; (dx^2 + dy^2) \\ &- \tfrac{4\,(1-\gamma)\,(z_2-z_1)}{\alpha^2\,(2-\gamma)^2\,\rho_0\, z_1^3} \left(1 - \tfrac{z}{z_1}\right)^{\frac{-8\,\gamma+8\,\epsilon+6}{7\,\gamma-4\,\epsilon-6}} \left(1 - \tfrac{z}{\alpha\, z_1}\right)^{\frac{4(5\,\gamma^2-2\,\gamma\,\epsilon-11\,\gamma+2\,\epsilon+6)}{(2-\gamma)(7\,\gamma-4\,\epsilon-6)}} \; dz^2 \end{aligned} \tag{32}$$

with $(1-\gamma)\,\rho_0\, z_1 > 0$.

Equation (29) is also readly integrable if $\epsilon \neq \gamma = 2$ or $\gamma \neq 2$ and $7\,\gamma = 4\,\epsilon + 6$, giving similar results. For $\gamma = 1$ or $\epsilon = \gamma = 2$, we must apply theorem (2). Therefore, the whole range of possibilities for constant γ and ϵ are easily covered.

In the literature, the perfect fluid $\epsilon = 0$ is most studied ([7] , [8], [9], [10], [11], [12], [2]-sec. 15.7.1), followed by the Einstein-Maxwell system of a charged infinite plane ([3], [13]), where $\gamma = 0$ and $\epsilon = -1$. The reader is invited to analyse in those works the effort one had to make solving Einstein equations for very specific values of γ and ϵ, whilst in our approach it is no more than a simple application of theorem 1, which gives general formulas just like in eq.(32).

We could have gone even further inserting a cosmological constant Λ from the beginning: just take $\rho+\Lambda$ and $p-\Lambda$ instead of ρ and p, respectively. As an example, the family of vacuum solutions with cosmological constant ([14,15]) is straightforwardly integrated from eq.(8) after such substitution is made and the "new" observables attained the values $\rho = p = \delta p = 0$.

4 Concluding remarks

In this work we have shown that any static plane-symmetric metric can be given, at least locally, as a simple functional of the matter content of space-time. In mathematics, it resembles the hard problem of characterizing the metric from its curvature tensor, which, in general, is impossible (See [16], sec. 4.5, for a short account of this problem in Riemannian geometry and the references therein). Contextualizing to general Relativity, due to the relevance of the generalized Birkhoff theorem ascribing a local form for the metric to vacuum solutions with certain symmetries [2], theorems 1 and 2 form a kind of "Birkhoff-type-theorem" with matter: *Given a certain class of symmetry assumed by* *the space-time and its matter content, decide if it is possible to find local coordinates for which the metric can be put as an explicit functional of the components of the energy-momentum tensor $T_{\mu\nu}$ such that both satisfy Einstein equations under the sole hypothesis of conservation $\nabla_\mu T^\mu_\nu = 0$, where the later corresponds to self-gravitating equations for the energy and momentum of the system.*

The results obtained so far purport to be valuable in the foundations of General Relativity. As a particular interest, grasping the relationship between the quantum Casimir effect and gravitation was the starting point of this work [17]. From this viewpoint, we have just set the ground where the problem may lay, and grasping it is a long, deep and uncertain journey. Investigations are under way.

References

[1] L. D. Landau , E. M. Lifshitz, : *Fluid Mechanics*, 2^{nd} ed. , Butterworth-Heinemann , 1987.

[2] H. Stephani, D. Kramer, M. Maccallum, C. Hoenselaers, E. Herlt: *Exact Solutions to Einstein's Field Equations*, Second Edition, Cambridge , 2003.

[3] P. A. Amundsen, Ø. Grøn: *General static plane-symmetric solutions of the Einstein-Maxwell equations*, Phys. Rev. D , Vol. 27, **8**, 1983.

[4] S. Weinberg: *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity*, Jonh Wiley & Sons, 1967.

[5] Ø. Grøn, S. Hervik: *Einstein's General Theory of Relativity*, Springer, 2007.

[6] A. D. Dolgov, I. B. Khriplovich: *Does a Static Solution Exist for a Gravitating Planar Wall?*, Gen. Rel. and Grav. , Vol. 21, **1**, 1989.

[7] A. H. Taub : *Isentropic hydrodynamics in plane symmetric space-times*, Phys. Rev. , vol. 103, **2**, 1956.

[8] R. Tabensky, A. H. Taub : *Plane Symetric Self-gravitating Fluids with Preassure Equal to Energy Density*, Commun. Math. Phys. Rev. , **29**, 1973.

[9] A. F. F. Teixeira, I. Wolk, M. M. Som : *Exact relativistic solution of disordered radiation with planar symmetry*, J. Phys. A , **10**, 1977.

[10] K. A. Bronnikov, M. A. Kovalchuk: *Properties of static fluid cylinders and plane layers in general relativity*, Gen. Rel. Grav. **11**, 1979.

[11] C. B. Collins : *Static relativistic perfect fluids with spherical, plane, or hyperbolic symmetry*, J. Math. Phys. **26** (9), September 1985.

[12] R. E. G. Saraví : *Static plane symmetric relativistic fluids and empty repelling singular boundaries*, Class. Quantum Grav. , **25**, 2008

[13] G. C. McVittie : *On Einstein's Unified Field Theory*, Proc. R. Soc. London A, **124**, 1929.

[14] J. Horský : *The gravitational field of a homogeneous plate with a non-zero cosmological constant*, Czech. J. Phys. B , **25**, 1975.

[15] J. Novotný, J. Horský : *On the plane gravitational condensor with the positive gravitational constant*, Czech. J. Phys. B , **24**, 1974.

[16] M. Berger: *A Panoramic View of Riemannian Geometry*, Springer, 2003.

[17] V. de Lorenci, L.G. Gomes, E. Moreira Jr: *Gravitational spectral shift caused by Casimir stresses*, Phys. Rev. D **84**, 2011.